



LAPIN YLIOPISTO
UNIVERSITY OF LAPLAND

University of Lapland



This is a self-archived version of an original article. This version usually differs somewhat from the publisher's final version, if the self-archived version is the accepted author manuscript.

Focus on recent, present and future Arctic and boreal productivity and biomass changes

Tømmervik, Hans; Forbes, Bruce C.

Published in:
Environmental research letters

DOI:
[10.1088/1748-9326/ab79e3](https://doi.org/10.1088/1748-9326/ab79e3)

E-pub ahead of print: 25.02.2020

Document Version
Peer reviewed version

Citation for published version (APA):
Tømmervik, H., & Forbes, B. C. (2020). Focus on recent, present and future Arctic and boreal productivity and biomass changes. *Environmental research letters*, 15(8), [080201]. <https://doi.org/10.1088/1748-9326/ab79e3>

Document License
CC BY

ACCEPTED MANUSCRIPT • OPEN ACCESS

Focus on recent, present and future Arctic and boreal productivity and biomass changes

To cite this article before publication: Hans Tømmervik *et al* 2020 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/ab79e3>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2020 The Author(s). Published by IOP Publishing Ltd.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

Focus on Recent, Present and Future Arctic and Boreal Productivity and Biomass Changes

Hans Tømmervik¹⁾ and Bruce C. Forbes²⁾

¹⁾Norwegian Institute for Nature Research, FRAM – High North Centre for Climate and the Environment, NO-9296 Tromsø, Norway,

²⁾Arctic Centre, University of Lapland, FI-96101 Rovaniemi, Finland

Abstract

The reduction of cold temperature constraints on photosynthesis in recent decades has led to extended growing seasons and increased plant productivity (greening) in significant parts of Polar, Arctic and Boreal regions, here called northern lands. However, most territories within these regions display stable productivity in recent years. Smaller portions of Arctic and Boreal regions show reduced productivity (browning). Summer drought and wildfires are the best documented drivers causing browning of continental areas. Yet factors like winter warming events dampening the greening effect of more maritime regions have remained elusive, least monitored and least understood. A Norway-US network project called ArcticBiomass was launched in 2013 to further reveal both positive and negative effects of climate change on biomass in Arctic and Boreal regions. This focus collection named *Focus on Recent, Present and Future Arctic and Boreal Productivity and Biomass Changes* includes 24 articles and is an important outcome of this work and addresses recent changes in phenology, biomass and productivity and the mechanisms. These mechanisms include former human interactions (legacies) and drivers that control such changes (both greening and browning), along with consequences for local, regional and global scale processes. We complete our synthesis by stressing remaining challenges and knowledge gaps, and

provide an outlook on future needs and research questions in the study of climate and human driven interactions in terrestrial Arctic and Boreal ecosystems.

1 Introduction

Ecosystem responses to Arctic warming have the potential to feedback either positively or negatively to the earth's climate system depending on latitude, changes in disturbance regime, vegetation distribution and productivity (McGuire et al. 2009).

The lower albedo of shrub and forest vegetation compared with tundra, for example, results in a positive feedback on temperature (Bala et al. 2007, Lorant et al. 2011).

Global warming presented as a warming rate, e.g. 0.2 °C/decade (ACIA 2004), does not reflect the fact that cold seasons are warming faster than the warm seasons, especially in the Arctic due to positive feedbacks (e.g. albedo-temperature feedback (Meredith et al. 2019)). The initiation, termination and performance of many biological processes, e.g. plant growth, are tied to threshold temperatures. The trend in timing of these thresholds, and cumulative air temperatures driving them, may have the effect of enhancing vegetation productivity.

Conversely, increased productivity of Arctic vegetation resulting from warmer temperatures tends to result in increased carbon dioxide (CO₂) uptake by net photosynthesis, providing a negative feedback to rising temperatures (Field et al. 2007; Serreze et al. 2000, Speed et al. 2010). As a result of this trend, between 1982 and 2011, Arctic tundra vegetation increased both in terms of peak productivity, greening and growing season length and this finding is supported by a wide range of field site measurements across the Arctic (Myneni et al. 1997, ACIA 2004; Serreze et al. 2000; Walker et al. 2006, Zu et al. 2013). Also, in coastal areas of Canada and Alaska (Epstein et al. 2004) and Arctic islands like Svalbard, there are trends of increased greening (Speed et al. 2010, Vickers et al. 2016), with some demonstrated linkages to

1
2
3
4 sea ice (Macias-Fauria et al. 2017). These dynamics include changes in the
5
6 composition and density of herbaceous vegetation (Epstein et al. 2004; Shaver et al.
7
8 2007), increased woody shrub encroachment in tundra areas (Sturm et al. 2001; Tape
9
10 et al. 2006, Tømmervik et al. 2009), increased height of in situ erect shrubs (Forbes et
11
12 al. 2010; Macias-Fauria et al. 2012; Bjorkman et al. 2018), changes in the depth of
13
14 seasonal thaw (Goulden et al. 1998; Kimball et al. 2006; Schuur et al. 2009; Loranty et
15
16 al. 2018), and associated changes in the energy regime (Chapin et al. 2005; Sturm et al.
17
18 2001). These insights are not obvious from measurements of warming rates alone, and
19
20 the relative importance of these competing feedbacks. Thus, the cumulative effect of
21
22 changing Arctic vegetation on the climate system and CO₂ fluxes, is still not very well
23
24 known, particularly in Svalbard (Speed et al. 2010; Johansen et al. 2012, Johansen &
25
26 Tømmervik 2014), parts of Alaska (references?), and Fennoscandia (Väisänen et al.
27
28 2014). Drought can modify these feedback effects, decoupling warming and
29
30 productivity as well as the balance of gross photosynthesis and plant respiration, which
31
32 varies substantially across plant functional types (Chapin et al. 1996; Goetz and Prince
33
34 1999).

35
36
37
38
39
40
41
42
43 The only effective way to map biomass and plant productivity in such large and remote
44
45 areas as Alaska and Svalbard, as well as in other territories of the Arctic, is using
46
47 remote sensing together with necessary *in situ* measurements and observations
48
49 (Myneni et al 1997, Karlsen et al. 2009, Beck et al. 2011).

50
51
52
53
54 The aim of our synthesis is to provide an overview of the studies in this focus issue
55
56 and to place their findings within the broader context of ecosystem– climate dynamics.

57
58
59 Our overview is organized into four sections that focus on studies that were primarily
60

relevant to: (1) Changes in the physical environment over high latitude regions and associated ecological and phenological changes in Arctic/Boreal vegetation, including vegetation-mediated responses and climate feedbacks; (2) Actual and potential biomass change influenced by (local) climate, natural disturbances, human impacts (e.g. resource extraction and legacies) and impacts on humans; (3) Transformation of open tundra vegetation to tall shrub tundra or forests, due to warming and other processes, influencing local and global climate, albedo and climate feedback mechanisms; and (4) Integration of *in situ* observations and manipulation experiments including remote sensing and other data sources to advance methodological approaches for measuring and monitoring.

We then discuss the key collective advances made in each of these areas by the studies within this focus issue, as well as opportunities for future research.

2 Northern lands (45° N)

The synthesis comprise studies conducted north of 45°N comprising both the Boreal and Arctic zones and in Fig. 1 and Table 1 we can see the areas that were covered by the different studies. In order to define both regions, the latest version of the MODIS International Geosphere-Biosphere Programme (IGBP) land cover map (Friedl et al 2010, WWW-MCD12Q1) and the Circumpolar Arctic Vegetation Map (CAVM) (Walker et al 2005, WWW-CAVM) is used. Arctic (8.16 million km²) is defined as the vegetated area north of 65°N, excluding agricultural land and forests, but including the tundra south of 65°N. Boreal region (17.86 million km²) is defined as the vegetated area between 45°N and 65°N, excluding agricultural land, tundra and nemoral forests.

3 Overview synthesis

3.1 *Changes in the physical environment over high latitude regions and associated ecological and phenological changes in Arctic/Boreal vegetation, including vegetation-mediated responses and climate feedbacks*

Changes and variation over time in the physical environment across northern high latitude regions have been demonstrated to be large. Malnes et al. (2017) report on the duration of snow season using snow cover area fraction data based on satellite data (daily 500 m standard snow product - MOD10A1 from MODIS) from the northernmost Norway. The first and last snow-free days for the study area were extracted from satellite data and compared to *in situ* met-station measurements. The start of the snow-free periods was up to 30 days later in spring 2000 and 2005 (DOY 135) compared with spring 2002 and 2006, which had an early start (DOY 105). The end of autumn/start of winter had a very late start in the years 2000 and 2007 (DOY 290) compared with 2009, whereas the first snow fall was 20 days earlier (270). In spring, the correlation between the first snow-free day mapped by MODIS data and snow data from 36 of 40 meteorological stations was highly significant ($p < 0.05$), with a of bias of less than 10 days for 34 of the stations. In autumn, 31 of 40 stations show highly significant ($p < 0.05$) correlation with MODIS data, and the bias was less than 10 days for 27 of the stations.

Park et al. (2016) analyzed satellite observations during 33 years in order to assess changes in growing season metrics (onset: SOS, end: EOS and length: LOS) and seasonal total gross primary productivity using the growing season summed NDVI

(GSSNDVI). They found that LOS had lengthened by 2.60 d dec^{-1} ($p < 0.05$) due to an earlier onset of SOS (-1.61 d dec^{-1} , $p < 0.05$) and a delayed EOS (0.67 d dec^{-1} , $p < 0.1$) the past three decades. The changes in growing season metrics were stronger in Eurasia (EA) and in boreal regions compared to North America (NA) and the arctic regions.

Reports on tundra greening are numerous and varied, and include increases in summer, spring, and autumn temperatures, as well as growing season length (Macias-Fauria et al. 2012; Zeng et al. 2013; Bhatt et al., 2017; Vickers et al., 2017). The paper by Vickers et al. (2016) made use of 30 years of a 1 km spatial resolution dataset (AVHRR) from 1986 to 2015 to calculate annual maximum NDVI over parts of Svalbard, and they found positive trends in maximum NDVI (+29%) and average summer temperature (+59%), which were significantly positively correlated with each other. On the same high arctic island, Karlsen et al. (2017) found a non-significant increase of 15% for the integrated MODIS NDVI defined as OP NDVI (onset to peak NDVI) for the period 2000-2014. At both local and regional scales, the latter NDVI-measure was found to predict biomass.

However, an increasing number of northern regions currently show declining productivity (browning). Such trends are evident both in North America and Eurasia, and factors assumed to contribute to this decline include recent reductions in summer moisture (cf. Verbyla 2015). Browning has also been attributed to mire and pond formation from increasing precipitation in some regions (Miles & Esau 2016, Lara et al. 2018), increasing moisture stress in other regions (Verbyla 2015, Sulla-Menashe et al. 2018), thermokarst development (Raynolds & Walker 2016), wildfire disturbance

(Chu et al. 2016), forest insect outbreaks (Bjerke et al. 2014, Sweet et al. 2015), increasing plant stress from winter warming and reduced snow protection (Bjerke et al. 2017, Treharne et al. 2018), fungal infestations and moose damage on young pine forests in the Nordic region (Marntell 2019), spring freeze damage after bud swelling (Chamberlain et al. 2019) and increasing industrial development (Hofgaard et al. 2010, Miles & Esau 2016) and finally general forest change, which may lead to temporal browning including browning due to logging of forests (White et al. 2017).

te Beest et al. (2016) found that when reindeer reduce shrub abundance/cover and height, summer albedo increases in both *Betula nana*-dominated heath vegetation and *Salix glauca*-dominated shrub tundra. Model results reveal associated lower net radiation, together with latent and sensible heat fluxes in heavily-grazed areas in all shrub-dominated vegetation types. Hence reindeer have a potential cooling effect on climate by increasing summer albedo and decreasing net radiation. Herbivory may have long-term consequences in warmed and ambient high Arctic tundra according to the study by Little et al (2017). They found that significantly more dead vascular plant material was found within warmed open-top chambers during a 12-year period compared to ambient plots, regardless of grazing history, but in contrast to many short-term experiments no difference in the amount of living material was found.

Belowground plant biomass allocation in tundra ecosystems and its relationship with temperature is of importance for modelling and analysis of climate change (Wang et al. 2016). They found that plant community biomass–temperature relationships were significantly different between above and belowground biomass. Tundra ecosystems

through altered litter input and distribution in the soil, as well as possible changes in root turnover.

The timing and duration of different pheno-phases within a plant's life cycle are critical for plant performance and growth. In the High Arctic, the start of many of these phenological phases is determined by the start of snowmelt, which can change in a changing climate (Semenchuk et al. 2016). In order to assess any change in these phases, Semenchuk et al. (2016) tested if snowmelt data control the timing and duration of phenological periods in Svalbard using a timing gradient from natural to experimentally altered snow depths. All pheno-phases followed irrespective of timing of occurrence, vegetative or reproductive nature, and three of the four phenological periods were fixed for most species such as the indigenous, hence the last was aperiodic. Semenchuk et al. (2016) concluded that periodic species like the dwarf shrubs *Dryas octopetala* and *Cassiope tetragona* and the grass species *Luzula arcuata* are likely to be limited in their ability to adapt to changing snowmelt dates, and thus may be disadvantaged compared to some of the invading species on Svalbard such as the herb *Rumex longifolius* the grass *Deschampsia cespitosa*. Gillespie et al (2016) studied the plant-pollinator interactions in High Arctic (Svalbard) and in their study they altered the timing of flowering phenology, using snow fences and open-top chambers. As expected, deep snow plots delay snow melt timing and this in turn delay the first and peak flowering dates of the plants, hence shortened the prefloration period overall. The OTCs, however, counteracted the delay in first and peak flowering to some extent. There was no effect of treatment on length of flowering season, although for all variables there were species-specific responses. The insect flower-visitor community was species poor, and although evidence of disruption to phenological

overlaps was not found, reduced insect-flower visitation rates to flowers in plots with deep snow may have occurred, due to limited observation methods.

3.2 Actual and potential biomass change and productivity influenced by (local) climate, natural disturbances, human impacts (e.g. resource extraction and legacies) and impacts on humans

Bjerke et al. (2015) detail the effects of above ground ice accumulation on meadow productivity by using ground observation and remotely sensed data (GIMMS NDVI3g data). Five contrasting snow season types were identified; snow-rich season with no soil frost or no ground-ice through low snow and considerable soil frost and ground-ice. Conditions of shallow snow depth and shallow soil frost or above-ground ice formation which may be a result of more frequent warming events, are rare at present but are predicted to become the dominant snow season type in Low Arctic and the Boreal lowlands. Agricultural productivity was lowest after winters with high accumulation of plant-damaging, hermetic above-ground ice formation Deep soil frost by itself did not reduce primary productivity. Lorant et al. (2016) quantified the distribution of vegetation productivity trends, wildfire, and near-surface soil carbon, according to vegetation type, across the continuous and discontinuous permafrost. Zones. They observed positive trends in vegetation productivity in areas of continuous permafrost, whereas areas underlain by discontinuous permafrost have proportionally less positive productivity trends.

3.3 Transformation of open tundra vegetation to shrub tundra or forests, due to warming and other processes, influencing local and global climate, albedo and climate feedback mechanisms.

In recent decades, woody shrubs have either increased in biomass/height and/or expanded into new areas throughout the Pan-Arctic tundra biome (Martin et al. 2017;

see also Forbes et al. 2010; Macias-Fauria et al. 2012). The same authors created a protocol for (a) identification of an operational suite of controls on shrub growth and expansion, and (b) characterization of an evidence base for controls on Arctic shrub growth and expansion. Evidence for a suite of 23 proximal controls that operate directly on shrub growth and expansion was found, while the evidence base was only focused on just four controls like air temperature, snow dynamics, soil moisture and herbivory. In particular, 65% of the evidence was generated within the warmest tundra climates (i.e. Low Arctic), while 24% of the evidence was from only one of 28 floristic sectors, indicating huge gaps (Martin et al. 2016) in the available evidence (Fig. 1) in), so not comprehensive enough to answer key questions concerning Pan-Arctic shrub change.

Significant expansion of shrubs has been observed within Arctic and Boreal regions during recent decades and in eastern sub-arctic Canada, where dendrochronological studies have demonstrated that the majority of shrub stands sampled were young, since the dominant stems were developed after 1990 (Paradis et al. 2016) which seem to be in accordance with Park et al. (2016). Stratified sampling of shrubs revealed that woody biomass was maximal within the lower canopy stratum, whereas foliar biomass tracked the development of the respective stands' vertical structure. Shrub height as a parameter explained snow depth, winter ground level temperature and summed freezing-degree days, while woody biomass best explained summer ground level temperature. Hence, shrub canopy structure will exert significant control on the abiotic environment in subarctic ecosystems (Paradis et al. 2016).

Shrub expansion is more likely to occur in areas with high soil moisture and nutrient availability, conditions typically found in sub-surface water channels called water tracks. Curasi et al. (2016) quantified the distribution of water tracks and their contribution CO₂ dynamics during the growing season for a Siberian tundra landscape using field measurements, satellite observations and meteorological data. They found that water tracks occupied 7.4% of the study area, and account for a slightly larger proportion of growing season carbon uptake relative to surrounding tundra and also larger relative to graminoids within the same water tracks. Water tracks are an

important component of this landscape and they will influence ecosystem structural and functional responses to climate, and is therefore of importance for modeling.

Increased wetness confounded Landsat-derived NDVI trends central Alaska North Slope region (Raynolds and Walker 2016) during 1985 to 2011. Regional trends showed decreases in NDVI for most vegetation types, but increases in tasseled-cap greenness, greatest for shrub dominated vegetation, and tasseled-cap wetness. This was consistent with thawing of polygon ice wedges. Increasing cover of water may be masking increases in vegetation when summarized using the more water-sensitive NDVI. This is also consistent with reduced NDVI due to melting of tundra as well as mire and pond formation from increasing precipitation in some regions of Northwest Siberia (Miles & Esau 2016), which show that care should be taken when relying solely on NDVI data (Raynolds and Walker 2016).

Disturbances can have particularly large effects on Arctic ecosystems when ecosystem structure and function are controlled by strong feedbacks between soil conditions, vegetation moisture, and sub-surface thermal regime (Cameron et al 2016). An example of such disturbance includes e.g. road construction and maintenance on vegetation structure and biomass along the Dempster Highway. Using field data and very high-resolution remotely sensed data, Cameron et al. (2016) found that increased shrub proliferation adjacent to the road was caused by greater soil moisture (see also Myers-Smith et al. 2015).




Figure 1. Map demonstrating regions of the Arctic for which there were evidence gaps during the period January 2012–January 2017 (Martin et al. 2017).

3.4 *Integration of in situ observations and manipulation experiments including remote sensing and other data sources to advance methodological approaches for measuring and monitoring.*

Buchhorn et al. (2016) showed that satellites provide the only practical source of data for estimating biomass of large and remote areas of the Arctic. Researchers have found that the normalized difference vegetation index (NDVI) correlates well with biomass sampled on the ground. However, errors in NDVI and biomass estimates due to bidirectional reflectance distribution function (BRDF) effects are not well reported. Finally, they found that studies that only sampling a narrow range of biomass and NDVI may produce equations that are more difficult to correct for BRDF effects.

Brazhnik and Shugart (2016) applied the new spatially-explicit gap-dynamics model SIBBORK towards a better understanding of how transition zones, namely treelines or forest lines, which are under-sampled and difficult to model, may change in the near future. They found that a 2 °C change in annual average air temperature will significantly alter the structure, composition, and productivity of boreal forest stands both at the northern treeline by 2040, and at the southern treeline by 2050.

Juutinen et al. (2017) assessed the spatial variation and seasonal dynamics of leaf-area index (LAI) linking ground observations and very-high-spatial resolution multispectral satellite images (e.g. Worldview-2). They illustrated how the short growing season, rapid development of the LAI, yearly climatic variation, and timing of satellite data should be accounted for in matching imagery and field verification data in the Arctic. Among the main plant functional types, graminoid LAI displayed the largest seasonal amplitudes and was the main cause of varying NDVI spatial patterning.

The utilization of earth observation data in vegetation monitoring is highly dependent on a long heritage of ground-based observations in the Arctic (Walker et al. 2016). Several products of the Conservation of Arctic Flora and Fauna are key to our current understanding (Shuchman et al. 2015, Christen T et al. 2013)) They have concluded that there is an urgent need for more consistent standards of plot-based observations and recommend improvements regarding the linkage between plot-based observations biodiversity studies and satellite-based observations of Arctic vegetation.

3.5 *New tools*

Bratsch et al (2017) assessed the ability of hyperspectral remote sensing data to estimate low arctic tundra biomass in Alaska. The main result was that the ability to identify unique biomass-spectra relationships within respective vegetation types using hyperspectral sensors was decreased during the peak of the growing season, since shrubs obscure lower-statured, bryophyte-dominated vegetation types. Hence, this study supports previous studies that shrubs control the spectral reflectance in Low Arctic communities (see also Forbes et al. 2010).

4 Further work and concluding remarks

Winter warming induced damage (arctic browning), particularly on dwarf-shrub vegetation (e.g. *Cassiope tetragona*) should be followed up via field monitoring and remote sensing (including UAV) in the future. Other vegetation changes in Arctic tundra and boreal regions, including the warming and thawing of permafrost (cf. Loranty et al. 2016, 2018), should be followed up. UAV based sensors and satellite sensors with very high spatial resolution (30-50 cm) may be efficient tools for detection of coastal erosion, human impacts (tourism), changes in tundra vegetation cover/stature, as well as permafrost thaw, and should be followed annually in the future. UAV can also be used as an upscaling tool. Another problem is that there exists huge gaps (Martin et al. 2017) in the Arctic and Boreal regions concerning long-term monitoring sites, including large areas that are not monitored at all (Fig. 1). These gaps currently afford us insufficient evidence of what is transpiring ecologically within these regions. However, a recent paper published by Virkkala et al (2019) in ERL provide detailed maps of potential new sampling locations in Arctic and the northern boreal region (the land north of the Arctic Circle (66,5° N), hence help prioritize future research efforts concerning environmental change in the north.

Acknowledgements

Financial support to this study was received from the Research Council of Norway through the projects ArcticBiomass, Svalbard Biomass and VanWhite (Research Council of Norway, contracts no. 227064, 270992 and 287402). BCF was also supported by Academy of Finland projects RISES (no. 256991) and HUMANOR (no. 251111; also JPI Climate no. 291581).

References (underlined articles belongs to the focus collection)

- Bhatt U et al 2017 Changing seasonality of Panarctic tundra vegetation in relationship to climatic variables. Environ. Res. Lett. 12 055003.
- Bjerke J W et al 2015 Impacts of snow season on ground-ice accumulation, soil frost and primary productivity in .a grassland of sub-Arctic Norway Environ. Res. Lett. 10 095007
- Bjorkman A D et al 2018 Changes in plant functional traits across a warming tundra biome. Nature **62** 57–62
- Bratsch S et al 2017 Relationships between hyperspectral data and components of vegetation biomass in Low Arctic tundra communities at Iivotuk, Alaska Environ. Res. Lett. 12 025003
- Brazhnik K and Shugart H H 2015 3D simulation of boreal forests: structure and dynamics in complex terrain and in a changing climate Environ. Res. Lett. 10 105006
- Buchhorn M et al 2016 Influence of BRDF on NDVI and biomass estimations of Alaska Arctic tundra Environ. Res. Lett. 11 125002
- Cameron E A, Lantz T C 2016 Drivers of tall shrub proliferation adjacent to the Dempster Highway, Northwest Territories, Canada Environ. Res. Lett. 11 04500
- Christensen T et al 2013. The Arctic Terrestrial Biodiversity Monitoring Plan. CAFF Monitoring Series Report Nr. 7. CAFF International Secretariat. Akureyri, Iceland. ISBN 978-9935-431-26-4
- Curasi S R et al 2016 Water track distribution and effects on carbon dioxide flux in an eastern Siberian upland tundra landscape Environ. Res. Lett. 11 045002
- DeFries R S et al. 1999 Combining satellite data and biogeochemical models to estimate global effects of human-induced land cover change on carbon emissions and primary productivity. *Global Biogeochem. Cy.* **13** 803-815
- Forbes B C, Fauria M M, Zetterberg P 2010 Russian Arctic warming and ‘greening’ are closely tracked by tundra shrub willows. *Glob. Change Biol.* **16** 1542–1554.
- Gillespie M A K et al 2016 High Arctic flowering phenology and plant–pollinator interactions in response to delayed snow melt and simulated warming Environ. Res. Lett. 11 115006

- Hofgaard *et al* 2010 Role of disturbed vegetation in mapping the boreal zone in northern Eurasia *Appl. Veg. Sci.* **13** 460–72
- Høgda K A, Tømmervik H and Karlsen S R 2013 Trends in the start of the growing season in Fennoscandia 1982–2011 *Remote Sens.* **5** 4304–4318
- Jepsen J U *et al* 2009 Monitoring the spatio-temporal dynamics of geometrid moth outbreaks in birch forest using MODIS-NDVI data. *Remote Sens. Environ.* **113** 1939–47
- Juutinen S *et al* 2017 Spatial variation and seasonal dynamics of leaf-area index in the arctic tundra-implications for linking ground observations and satellite images *Environ. Res. Lett.* **12** 095002
- Karlsen S R *et al* 2009 Growing-season trends in Fennoscandia 1982–2006, determined from satellite and phenology data. *Clim. Res.* **39** 275–86
- Karlsen, S R, Anderson H B, van der Wal R, Hansen B B 2018 A new NDVI measure that overcomes data sparsity in cloud-covered regions predicts annual variation in ground-based estimates of high arctic plant productivity *Environ. Res. Lett.* **13** 025011
- Kuuluvainen T *et al* 2017 North Fennoscandian mountain forests: History, composition, disturbance dynamics and the unpredictable future *Forest Ecol. Manag.* **385** 140–49
- Little C J *et al* 2017 Short-term herbivory has long-term consequences in warmed and ambient high Arctic tundra *Environ. Res. Lett.* **12** 025001
- Loranty M M *et al* 2016 Spatial variation in vegetation productivity trends, fire disturbance, and soil carbon across arctic-boreal permafrost ecosystems *Environ. Res. Lett.* **11** 095008
- Loranty M M *et al* 2018 Changing ecosystem influences on soil thermal regimes in northern high-latitude permafrost regions. *Biogeosciences* **15** 5287–5313,
- Macias-Fauria M, Forbes, B C, Zetterberg P, Kumpula T 2012 Eurasian Arctic greening reveals teleconnections and the potential for structurally novel ecosystems. *Nat. Clim. Change* **2** 613–618.
- Macias-Fauria M, Karlsen S R, Forbes B C 2017 Disentangling the coupling between sea ice and tundra productivity in Svalbard. *Scientific Reports* **7** 8586
- Malnes E *et al* 2016 Snow season variability in a boreal-Arctic transition area monitored by MODIS data *Environ. Res. Lett.* **11** 125005

- Martin A C et al 2017 Shrub growth and expansion in the Arctic tundra: an assessment of controlling factors using an evidence-based approach. *Environ. Res. Lett.*, 12, 085007
- Meredith M et al 2019 Chapter 3, Polar Regions, In: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate. UNEP/WMO.
- Miles V V, Esau I 2016 Spatial heterogeneity of greening and browning between and within bioclimatic zones in northern West Siberia *Environ. Res. Lett.* 11 115002
- Normand S et al 2017 Legacies of Historical Human Activities in Arctic Woody Plant Dynamics. *Annu. Rev. Env. Res.* 42 541–67
- Paradis M et al 2016 Greater effect of increasing shrub height on winter versus summer soil temperature *Environ. Res. Lett.* 11 085005
- Park et al 2016 Changes in growing season duration and productivity of northern vegetation inferred from long-term remote sensing data *Environ. Res. Lett.* 11 084001
- Raynolds M K, Walker D A 2016 Increased wetness confounds Landsat-derived NDVI trends in the central Alaska North Slope region, 1985–2011 *Environ. Res. Lett.* 11 085004
- Semenchuk P R et al 2016 High Arctic plant phenology is determined by snowmelt patterns but duration of phenological periods is fixed: an example of periodicity *Environ. Res. Lett.* 11 125006
- Shuchman R A, Jenkins L, Whitley M A 2015. Arctic Land Cover Change Initiative: MODIS Satellite Data. CAFF Monitoring Series No. 17. Conservation of Arctic Flora and Fauna, Akureyri, Iceland. ISBN: 978-9935-431-49-3.
- te Beest M et al Reindeer grazing increases summer albedo by reducing shrub abundance in Arctic tundra 2016 *Environ. Res. Lett.* 11 125013
- Tucker C J, Sellers P J 1986 Satellite remote sensing of primary production. *Int. J. Remote Sens.* 7 1395–1416.
- Verbyla D 2015 Remote sensing of interannual boreal forest NDVI in relation to climatic conditions in interior Alaska *Environ. Res. Lett.* 10 125016
- Vickers H et al 2016 Changes in greening in the high Arctic: insights from a 30 year AVHRR max NDVI dataset for Svalbard *Environ. Res. Lett.* 11 105004
- Virkkala et al 2019 Identifying multidisciplinary research gaps across Arctic terrestrial

- 1
2
3
4 gradients. *Environ. Res. Lett.* 14 124061
5
6 Väisänen M *et al* 2014 Consequences of warming on tundra carbon balance
7 determined by reindeer grazing history. *Nat. Clim. Change* 4 384–388
8
9 Walker D A *et al* 2012 Environment, vegetation and greenness (NDVI) along the
10 North America and Eurasia Arctic transects. *Environ. Res. Lett.* 7 015504
11
12 Walker D A *et al* 2012 Circumpolar Arctic vegetation: a hierarchic review and
13 roadmap toward an internationally consistent approach to survey, archive and
14 classify tundra plot data. *Environ. Res. Lett.* 11 055005
15
16 Wang P *et al* 2016 Belowground plant biomass allocation in tundra ecosystems and its
17 relationship with temperature *Environ. Res. Lett.* 11 055003
18
19 White J C *et al* 2017 Nationwide annual characterization of 25 years of forest
20 disturbance and recovery for Canada using Landsat time series. *Remote Sens.*
21 *Environ.* 194 303–321
22
23 Williams M *et al* 1997 Predicting gross primary productivity in terrestrial ecosystems
24 *Ecol. Appl.* 7 882–94
25
26 Xu L *et al* 2013 Temperature and vegetation seasonality diminishment over northern
27 lands *Nat. Clim. Change* 3 581–86
28
29 Zeng H, Jia G S, Forbes B C 2013 Response of phenological shifts to climate and
30 anthropogenic factors as detected from multi-satellite data. *Environ. Res. Lett.* 8
31 035036. doi:10.1088/1748-9326/8/3/035036.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60